

Solving Quadratic Equations by Completing the Square

A quadratic equation is an equation in the form of $ax^2 + bx + c = 0$ where **a**, **b**, and **c** are constants and $a \neq 0$. One of the methods that can be used to solve a quadratic equation is completing the square.

- Step 1:** Isolate the constant if it is not isolated.
- Step 2:** If the coefficient of x^2 is not 1, divide both side of the equation by the coefficient of x^2 .
- Step 3:** Divide the coefficient of x by 2, and square the result.
- Step 4:** Add the result from step 3 to both sides of the equation, and simplify.
- Step 5:** Factor the variable side of the equation.
(Hint: In order to have a complete square, both factors must be the same.)
- Step 6:** Take the square root of both sides of the equation.
Note: Place the \pm sign in front of square root of **constant**.
- Step 7:** Isolate x .

Example 1: $x^2 - 6x + 5 = 0$

Step 1: $x^2 - 6x = -5$

Step 2: skip

Step 3: $\frac{-6}{2} = -3$

$$(-3)^2 = 9$$

Step 4: $x^2 - 6x + 9 = -5 + 9$

$$x^2 - 6x + 9 = 4$$

Step 5: $(x - 3)(x - 3) = 4$

$$(x - 3)^2 = 4$$

Hint: -3 is half of the coefficient of x .

Step 6: $\sqrt{(x - 3)^2} = \pm \sqrt{4}$

$$x - 3 = \pm 2$$

Step 7: $x - 3 = -2$ or $x - 3 = 2$

$$x = 1$$
 or $x = 5$

Example 2: $3x^2 + 6x - 72 = 0$

Step 1: $3x^2 + 6x = 72$

Step 2: $\frac{3x^2}{3} + \frac{6x}{3} = \frac{72}{3}$

$$x^2 + 2x = 24$$

Step 3: $\frac{2}{2} = 1$

$$(1)^2 = 1$$

Step 4: $x^2 + 2x + 1 = 24 + 1$

$$x^2 + 2x + 1 = 25$$

Step 5: $(x + 1)(x + 1) = 25$

$$(x + 1)^2 = 25$$

Hint: 1 is half of 2 (the coefficient of x).

Step 6: $\sqrt{(x + 1)^2} = \pm \sqrt{25}$

$$x + 1 = \pm 5$$

Step 7: $x + 1 = -5$ or $x + 1 = 5$

$$x = -6$$
 or $x = 4$

Example 3: $x^2 + 6x + 10 = 0$

Step 1: $x^2 + 6x = -10$

Step 2: skip

Step 3: $\frac{6}{2} = 3$

$$\left(\frac{6}{2}\right)^2 = 9$$

Step 4: $x^2 + 6x + 9 = -10 + 9$

$$x^2 + 6x + 9 = -1$$

Step 5: $(x + 3)(x + 3) = -1$

$$(x + 3)^2 = -1$$

Hint: 3 is half of the coefficient of x.

Step 6: $\sqrt{(x + 3)^2} = \pm \sqrt{-1}$

$$x + 3 = \pm i$$

(Note: $\sqrt{-1} = i$)

Step 7: $x + 3 = -i$ or $x + 3 = i$

$$x = -3 - i$$
 or $x = -3 + i$

Example 4: $2x^2 + 5x - \frac{1}{2} = 0$

Step 1: $2x^2 + 5x = \frac{1}{2}$

Step 2: $\frac{2}{2}x^2 + \frac{5}{2}x = \frac{1/2}{2}$

$$x^2 + \frac{5}{2}x = \frac{1}{4}$$

Step 3: $\frac{5/2}{2} = \left(\frac{5}{2}\right)\left(\frac{1}{2}\right) = \frac{5}{4}$

$$\left(\frac{5}{2}\right)^2 = \frac{25}{16}$$

Step 4: $x^2 + \frac{5}{2}x + \frac{25}{16} = \frac{1}{4} + \frac{25}{16}$

$$x^2 + \frac{5}{2}x + \frac{25}{16} = \frac{29}{16}$$

Step 5: $\left(x + \frac{5}{4}\right)\left(x + \frac{5}{4}\right) = \frac{29}{16}$

$$\left(x + \frac{5}{4}\right)^2 = \frac{29}{16}$$

Hint: $\frac{5}{4}$ is half of $\frac{5}{2}$ (the coefficient of x.)

Step 6: $\sqrt{\left(x + \frac{5}{4}\right)^2} = \pm \sqrt{\frac{29}{16}}$

$$x + \frac{5}{4} = \pm \frac{\sqrt{29}}{4}$$

Step 7: $x + \frac{5}{4} = -\frac{\sqrt{29}}{4}$ or $x + \frac{5}{4} = \frac{\sqrt{29}}{4}$

$$x = \frac{-5 - \sqrt{29}}{4}$$
 or $x = \frac{-5 + \sqrt{29}}{4}$