

# Graphing Rational Functions

$$f(x) = \frac{N(x)}{D(x)} \text{ where } N \text{ and } D \text{ are polynomials with no common factors.}$$

I. **V.A.:** Find the **vertical asymptote(s)** by setting the denominator equal to zero:  $D(x) = 0$

II. **H.A.:** Find the **horizontal asymptote** by using the following rules:

$$f(x) = \frac{a_n x^n + a_{n-1} x^{n-1} + \dots}{b_m x^m + a_{m-1} x^{m-1} + \dots} \quad \text{The degree of the numerator is } n. \text{ The degree of denominator is } m.$$

a) If  $n < m$ , then  $y=0$  (x-axis) is the horizontal asymptote

b) If  $n=m$ , then  $y = \frac{a_n}{b_m}$  is the horizontal asymptote.

c) If  $n > m$ , then there is no horizontal asymptote

**Exception:** If  $n > m$  by 1, then the graph has a **slant asymptote (S.A.)**

III. Find the **x-intercepts** by setting the numerator equal to zero and solve.  $N(x) = 0$

IV. Find the **y-intercepts** by substituting zero for x in the function,  $f(0)$

V. Make a table of values between and beyond each x-intercept and vertical asymptote; *more values provide a more accurate graph.*

VI. Sketch the graph by using the information obtained.

**. NOTE: Remember that the graph never crosses the vertical asymptotes!!!!**

**EXAMPLE 1:**      *Graph the function -*  $f(x) = \frac{3x^2}{x^2 - 4}$

I. **V.A.:** Set the denominator equal to zero and solve.

$$\begin{aligned} x^2 - 4 &= 0 \\ x^2 &= 4 \\ x &= \pm 2 \end{aligned}$$

*Vertical asymptotes at*  
 $x = 2 \text{ \& } x = -2$

II. **H.A.:** The degree of the numerator and denominator are both equal to 2. So the horizontal asymptote is  $y = \frac{3}{1}$ .

*Horizontal asymptote at*  $y=3$

III. **x-intercept(s):** Set the numerator equal to zero and solve.

$$\begin{aligned} 3x^2 &= 0 \\ x^2 &= 0 \\ x &= 0 \end{aligned} \quad \text{Intercept at } x = 0$$


IV. **y-intercept(s):** Substitute zero for x in the function and solve.

$$\begin{aligned} y &= \frac{3(0)^2}{(0)^2 - 4} = -\frac{0}{4} \\ y &= 0 \end{aligned} \quad \text{Intercept at } y = 0$$

V. **Table:** Make a table of values near each vertical asymptote and x-intercept and plot the values

$x$	$f(x) = \frac{3x^2}{x^2 - 4}$
-4	4
-3	$\frac{27}{5}$
-1	-1
1	-1
3	$\frac{27}{5}$
4	$\frac{4}{4}$

VI. **Graph:** Sketch the graph by using the information obtained.



**EXAMPLE 2:** Graph the function -  $f(x) = \frac{x+3}{x^2-9}$

$$f(x) = \frac{x+3}{x^2-9} = \frac{\cancel{x+3}}{(\cancel{x+3})(x-3)} = \frac{1}{x-3}$$

VI. **Graph:**

I. **V.A.:**  $x - 3 = 0$   $x = 3$


II. **H.A.:**  $n < m$  so  $y = 0$

III. **x-intercept:**  $1 \neq 0$ , so there are none

IV. **y-intercept:**  $\frac{1}{0-3} = -\frac{1}{3}$

V. **Table:**

$x$	$f(x) = \frac{x+3}{x^2-9}$
2	-1
4	1



**EXAMPLE 3:**      *Graph the function -  $f(x) = \frac{x+3}{x^2+9}$*

I. **V.A.:**                       $x^2 + 9 \neq 0$   
*The function is always positive, so there are no vertical asymptotes*

II. **H.A.:**                       $n < m$  so  $y = 0$


III. **x-intercept:**             $x + 3 = 0$      $x = -3$

IV. **y-intercept:**             $\frac{0+3}{0^2+9} = \frac{1}{3}$

V. **Table:**

$x$	-5	-2	1	3
$f(x) = \frac{x+3}{x^2+9}$	$-\frac{1}{17}$	$\frac{1}{13}$	$\frac{2}{5}$	$\frac{1}{3}$

VI. **Graph:**



**EXAMPLE 4:**      *Graph the function -  $f(x) = \frac{x^2+1}{x-1}$*

I. **V.A.:**                       $x - 1 = 0$      $x = 1$

VI. **Graph:**

II. **H.A.:**                       $n > m$  by 1, therefore there are no horizontal asymptotes, but there is a slant asymptote; the equation can be found by division

$$f(x) = \frac{x^2+1}{x-1} = \boxed{x+1} + \frac{2}{x-1}$$

Equation of Slant Asymptote  
 $y = x+1$

III. **x-intercept:**             $x^2 + 1 = 0$      $x^2 \neq -1$   
*so there are none*

IV. **y-intercept:**             $\frac{0^2+1}{0-1} = -1$

V. **Table:**

$x$	$-\frac{1}{2}$	2
$f(x) = \frac{x^2+1}{x-1}$	$-\frac{5}{6}$	5

