

Vector Dot Product

Consider a vector \mathbf{A} (we use **boldface** to denote a vector) with rectangular coordinates A_x and A_y . We can write \mathbf{A} as follows:

$$\mathbf{A} = A_x\mathbf{i} + A_y\mathbf{j}$$

where \mathbf{i} is a unit vector in the x direction and \mathbf{j} is a unit vector in the y direction.

If we have two vectors \mathbf{A} and \mathbf{B} , defined as above, then we define the *dot product* of \mathbf{A} and \mathbf{B} , written $\mathbf{A} \cdot \mathbf{B}$ (read “A dot B”) as follows:

$$\mathbf{A} \cdot \mathbf{B} = A_xB_x + A_yB_y \quad (1)$$

Notice that the dot product is a *scalar*; it is the sum of one or more scalars. It may be referred to as the *scalar product*.

The following websites may also be helpful:

<http://www.falstad.com/dotproduct/>

<http://betterexplained.com/articles/vector-calculus-understanding-the-dot-product/>

Simple EXAMPLE:

Suppose $\mathbf{A} = 3\mathbf{i} + 5\mathbf{j}$ and $\mathbf{B} = 4\mathbf{i} - 2\mathbf{j}$

STEP 1: Identify the components of each vector

$$A_x = 3 \quad A_y = 5 \quad B_x = 4 \quad B_y = -2$$

STEP 2: Use equation (1) to compute the dot product

$$\mathbf{A} \cdot \mathbf{B} = A_xB_x + A_yB_y = (3)(4) + (5)(-2) = 12 - 10 = 2$$

More Complicated EXAMPLE:

The rules are the same for three-dimensional vectors.

Suppose $\mathbf{A} = 3\mathbf{i} + 5\mathbf{j} + 8\mathbf{k}$ and $\mathbf{B} = 4\mathbf{i} - 2\mathbf{j} + \mathbf{k}$. Compute $\mathbf{A} \cdot \mathbf{B}$.

STEP 1: Identify the components of each vector

$$A_x = 3 \quad A_y = 5 \quad A_z = 8 \quad B_x = 4 \quad B_y = -2 \quad B_z = 1$$

STEP 2: Use equation (1) to compute the dot product

$$\mathbf{A} \cdot \mathbf{B} = A_x B_x + A_y B_y + A_z B_z = (3)(4) + (5)(-2) + (8)(1) = 10$$

Alternate Formula

It happens that the dot product can also be calculated as follows:

$$\mathbf{A} \cdot \mathbf{B} = |\mathbf{A}| |\mathbf{B}| \cos \theta, \quad (2)$$

where $|\mathbf{A}|$ is the magnitude of \mathbf{A} , $|\mathbf{B}|$ is the magnitude of \mathbf{B} , and θ is the angle between them.

This formula may be helpful when you are given the magnitudes and angles to begin with.

EXAMPLE:

Suppose \mathbf{A} is a vector of magnitude 6 at an angle of 60° . Suppose \mathbf{B} is a vector of magnitude 4 at an angle of -25° . Compute $\mathbf{A} \cdot \mathbf{B}$.

STEP 1: Compute the angle θ between the vectors

$$\theta = 60^\circ - (-25^\circ) = 85^\circ$$

STEP 2: Compute the cosine of the angle θ

$$\cos \theta = \cos(85^\circ) \sim 0.0872 \text{ (rounding off)}$$

STEP 3: Use equation (2) to compute the dot product

$$\mathbf{A} \cdot \mathbf{B} = |\mathbf{A}| |\mathbf{B}| \cos \theta = (6)(4)(0.0872) \sim 2.09$$

Interpretation

The dot product of \mathbf{A} and \mathbf{B} can be thought of as the product of one vector times the component of the second that lies parallel to the first.

Note that $\mathbf{A} \cdot \mathbf{B}$ ranges from 0 when \mathbf{A} and \mathbf{B} are perpendicular to $|\mathbf{A}| |\mathbf{B}|$ when \mathbf{A} and \mathbf{B} are parallel.